

Available online at www.sciencedirect.com



Procedia Engineering 00 (2015) 000–000



www.elsevier.com/locate/procedia

24th International Meshing Roundtable (IMR24) On Indecomposable Polyhedra and The Number of Steiner Points

Nadja Goerigk, Hang Si

Weierstrass Institute, Mohrenstr. 39, 10117 Berlin, Germany

Abstract

The existence of *indecomposable polyhedra*, that is, the interior of every such polyhedron cannot be decomposed into a set of tetrahedra whose vertices are all of the given polyhedron, is well-known. However, the geometry and combinatorial structure of such polyhedra are much less studied. In this article, we investigate the structure of some well-known examples, the so-called *Schönhardt* polyhedron [10] and the Bagemihl's generalization of it [1], which will be called *Bagemihl's polyhedra*. We provide a construction of an additional point, so-called *Steiner point*, which can be used to decompose the Schönhardt and the Bagemihl's polyhedra. We then provide a construction of a larger class of three-dimensional indecomposable polyhedra which often appear in grid generation problems. We show that such polyhedra have the same combinatorial structure as the Schönhardt's and Bagemihl's polyhedra, but they may need more than one Steiner point to be decomposed. Given such a polyhedron with $n \ge 6$ vertices, we show that it can be decomposed by adding at most $\left\lceil \frac{n-5}{2} \right\rceil$ interior Steiner points. We also show that this number is optimal in the worst case.

© 2015 The Authors. Published by Elsevier Ltd.

Peer-review under responsibility of organizing committee of the 24th International Meshing Roundtable (IMR24).

Keywords: Indecomposable polyhedron, Steiner point, Schönhardt polyhedron; Bagemihl's polyhedron, tetrahedralization

1. Introduction

The existence of three-dimensional (non-convex) polyhedron whose interior cannot be decomposed into a set of non-overlapping tetrahedra without new vertices has long been observed [7]. In 1928, Schönhardt provided the simplest example, which is a twisted triangular prism with six vertices. It is now well-known as the *Schönhardt polyhedron* [10]. Later, further such non-convex, non-tetrahedralizable polyhedron with an arbitrary number of vertices have been presented, see [1,3,8]. Among them, Bagemihl's construction [1] is a direct generalization from the Schönhardt's construction.

The existence of indecomposable polyhedra is a major difficulty in many geometric and combinatorial problems. For example, Below shows that the complexity of finding minimal or maximal subdivisions of three-dimensional polyhedra are NP-hard [2]. Rupert and Seidel show that to determine whether a given three-dimensional polyhedron can be decomposed or not is NP-complete [9]. The Schönhardt polyhedron appears as an important example in the

1877-7058 © 2015 The Authors. Published by Elsevier Ltd.

Peer-review under responsibility of organizing committee of the 24th International Meshing Roundtable (IMR24).

E-mail address: Nadja Goerigk, Nadja.Goerigk@wias-berlin.de, Hang Si, Hang.Si@wias-berlin.de

study of the flip-graph of all triangulations of a given point set [4]. In 3d tetrahedral mesh generation, indecomposable polyhedra are the main obstacles in the design and proof of several key algorithms, such as the recovery of a non-existing edge and the deletion of an existing vertex, see e.g. [5,6,11,12].

When tetrahedralizing an indecomposable polyhedron, it is necessary to add additional points, the so-called *Steiner* points ¹. It is easy to show that the Schönhardt polyhedron needs at most one Steiner point. However, Chazelle constructed a polyhedron that may need as many as $\Omega(n^2)$ Steiner points. We can treat the Schönhardt polyhedron and Chazelle's polyhedron as the two extreme cases for the number of Steiner points. Surprisingly enough, very few works about the number of Steiner points between O(1) and $O(n^2)$ is known. The question we are going to investigate in this article is,

Given a 3d indecomposable polyhedron, how many Steiner points are necessary to decompose it?

In order to answer this question, it is necessary to understand the geometry and the combinatorial structure of the given polyhedron.

In this paper, we first study this question for the class of polyhedra constructed by Bagemihl [1], which is a generalisation of the Schönhardt polyhedron. Hereafter we will call them *Bagemihl's polyhedra* (described in Section 2.1). A Bagemihl's polyhedron can have arbitrary number of vertices. Due to its structural properties, we show that a Bagemihl's polyhedron needs only one interior Steiner point to be decomposed. Our proof is based on a construction of a Steiner point in the interior of a given Bagemihl's polyhedron, and we proof that all boundary faces of this polyhedron are visible by this Steiner point (in Section 2.2). Furthermore, we show that our constructed Steiner point is also valid for a class of *generalized Bagemihl's polyhedra* which can be obtained by relaxing the symmetric and height requirements in Bagemihl's construction (in Section 2.3).

Next, in Section 3, we first extends Bagemihl's Theorem to show that there exists a larger class of indecomposable polyhedra (in Section 3.1). We then give a general construction of such polyhedra with $n \ge 6$ vertices, denoted as σ_n . It is worth mentioning, that the polyhedra from our construction are commonly encountered during the process of tetrahedral mesh generation. We show that our constructed polyhedra have the same combinatorial structure as the Schönhardt's and Bagemihl's polyhedra, but they may need more than one interior Steiner point to be decomposed (in Section 3.2). We then prove the following main result regarding the number of Steiner points for our constructed polyhedra (in Section 3.3):

Given a polyhedron σ_n that satisfies our construction, where n is the number of vertices of σ_n , it needs at $most \left[\frac{n-5}{2}\right]$ interior Steiner points to be decomposed.

Our proof of this result is based on a construction of Steiner points in the interior of such polyhedra so that it can be decomposed into a set of tetrahedra with the help of these Steiner points. This construction also provides hints to design efficient algorithms to tetrahedralise such polyhedron.

Finally, some closing remarks and open questions are given in Section 4.

2. Bagemihl's Polyhedra and a Construction of a Steiner Point

In the Paper "On Indecomposable Polyhedra" by F. Bagemihl ([1]) he proves the following theorem.

Theorem 1 ([1]). *If n is an integer not less than* 6*, then there exists a polyhedron,* π_n *, with n vertices and the following properties:*

- (1) π_n is simple and every one of its faces is a triangle.
- (II) If τ is a tetrahedron, each of whose vertices is a vertex of π_n , then not every interior point of τ is an interior point of π_n .

¹ There exist several types of Steiner points, named after Jakob Steiner (1796 – 1863), a Swiss mathematician who worked primarily in geometry, in the literatures, like the Steiner points in the Steiner tree problem, see e.g. http://en.wikipedia.org/wiki/Steiner_tree_problem, and the Steiner point in a triangle, see e.g. http://en.wikipedia.org/wiki/Steiner_point_(triangle). The Steiner points in this article are constructed for decomposing Schönhardt or other indecomposable polyhedra. They are different to the previous ones.

- (III) Every open segment whose endpoints are vertices of π_n , but which is not an edge of π_n , lies wholly exterior to π_n .
- (IV) Every triangle whose sides are edges of π_n is a face of π_n .

Comments. In the original version of Theorem 1 [1], Bagemihl only stated the first three properties, (I), (II), and (III). However, he gave a construction of a class of polyhedra which also fulfills the property (IV). Note that by including the property (IV), we might decrease the size of the original class of polyhedra. But this is out of the scope of this article. With the included property (IV), the property (II) becomes redundant. It is followed together from (I), (III) and (IV).

Note that the property (II) indicates that π_n is indecomposable, since no tetrahedron τ whose vertices are from π_n in the interior of π_n exists. The key fact is that τ must contain at least one open segment of π_n . Suppose the four edges of τ are all not open segments, then, by (IV), the boundary of π_n must form a tetrahedron. Since π_n is simple by (I), we can conclude that π_n is a tetrahedron. But this is contradict to the assumption $n \ge 6$. Then by (III), τ does not lie the interior of π_n .

Bagemihl provides a construction of a class of polyhedra that satisfy Theorem 1, which will be called *Bagemihl's* polyhedra.

In this section, we will first review the construction of Bagemihl's polyhedra. Our goal is to show that a Bagemihl's polyhedron needs only one Steiner point to be decomposed. For this purpose, we first give a construction of a Steiner point in a given Bagemihl's polyhedron. We then prove it is valid for the decomposition. We further prove that our Steiner point is also valid for a variation of Bagemihl's polyhedra by relaxing the symmetry and edge length requirements in the original Bagemihl's construction.

2.1. Description of Bagemihl's Polyhedra

Bagemihl's construction starts with the Schönhardt polyhedron π_6 which we will describe first. Take an equilateral triangle with edge lengths 1 and vertices A_1, B_1, C_1 . Take a copy of it and lift it up orthogonally to the height h = 1 and rotate it around the axis connecting the centers of the top and bottom triangle by an angle of $\vartheta = 30^\circ$. Call the so obtained vertices in the top triangle A_2, B_2, C_2 , respectively. By connecting the vertices as shown in Figure 1 we obtain the polyhedron π_6 .



Fig. 1. The Schönhardt polyhedron, which is the Bagemihl's polyhedron π_6 . A side view (left) and a top view (right) are shown.

For the case n > 6, Bagemihl adds an open circular arc $\widehat{A_1A_2}$ connecting A_1 and A_2 in the interior of π_6 . The radius of this arc is chosen to be large enough such that every point of $\widehat{A_1A_2}$ is on the same side of the plane $C_1A_2C_2$ as A_1 , and on the same side of the plane $B_1A_1B_2$ as A_2 . On the arc $\widehat{A_1A_2}$ one can choose k = n - 6 distinct points, D_1, D_2, \ldots, D_k , in the order $A_1D_1D_2 \ldots D_{k-1}D_kA_2$ and add the edges $A_1D_1, D_1D_2, \ldots, D_{k-1}D_k, D_kA_2$ connecting the vertices. An example of a π_9 is shown in Figure 2.



Fig. 2. A Bagemihl's polyhedron π_9 . A side view (left) and a top view (right) are shown.

2.2. Construction of a Steiner Point

Given a Bagemihl's polyhedron, π_n with $n \ge 6$ vertices, it is clear that at least one Steiner point is needed. But it is not obvious how many Steiner points are necessary. We want to show that one Steiner point is already sufficient. In this section, we give a construction of such a Steiner point.

We first consider π_6 , which is just the Schönhardt polyhedron. The region in which we can place a Steiner point is the intersection of the eight halfspaces defined by the boundary triangles of the Schönhardt polyhedron, see Figure 3.



Fig. 3. The (open) valid domain for placing Steiner points inside the Schönhardt polyhedron. A side view (left) and a top view (right) are shown.

Placing the points D_1, \ldots, D_k , where k = n - 6, in the interior of π_6 just as described by Bagemihl, the region for a Steiner point *S* in π_n is getting smaller. We want to show that it is not empty for all $k \in \mathbb{N}_{\geq 0}$ and all choices of additional points D_i . For this purpose we will show that it is always possible to place a Steiner point in the interior that has the required visibility properties.

We first determine the domain in which the valid arcs as described by Bagemihl can live. First, every arc has to lie inside π_6 , so it is restricted by the halfspaces limited by the faces $A_1A_2C_1$ and $A_1A_2B_2$, respectively. Since every point of an open arc $\widehat{A_1A_2}$ has to be on the same side of the plane $C_1A_2C_2$ as A_1 , and on the same side of the plane $B_1A_1B_2$ as A_2 [1, p. 413], it is restricted by these two faces of π_6 as well. So, this domain is the intersection of four half spaces, which is a tetrahedron (see Figure 4), denoted as T, with vertices $A_1A_2G_1G_2$, where G_1 and G_2 are defined by

$$G_1 := \operatorname{plane}_{C_1 A_2 C_2} \cap \operatorname{line}_{A_1 B_2}$$
$$G_2 := \operatorname{plane}_{B_1 A_1 B_2} \cap \operatorname{line}_{A_2 C_1}.$$

Note that the points G_1 and G_2 are on the boundary of π_6 : Since the polyhedron doesn't have interior edges, the points A_1 and B_2 will lie on different sides of the plane through $C_1A_2C_2$, so G_1 has to be on the edge A_1B_2 itself. Analogously G_2 has to be on the edge A_2C_1 .

With the help of this tetrahedron T, we can determine the locations for the intersections of the tangent lines of all possible arcs at A_1 and A_2 . Given a possible arc $\widehat{A_1A_2}$ (as defined by Baghmil) in the tetrahedron T, define the intersection point of the tangent lines through A_1 and A_2 to the arc $\widehat{A_1A_2}$ as P (refer to Figure 4). P must lie in the closure of T. If the radius of the circle (containing the arc) is getting larger, the distance between P and the line through A_1 and A_2 will decrease.

It is enough to consider the extremal case, i.e. the case when the radius of the circle is the smallest possible.

Consider the points G_1 and G_2 . By definition, they can see all vertices of π_n or are coplanar with a face containing them. The open segment G_1G_2 is obtained by first intersecting the two planes through $C_1A_2C_2$ and $B_1A_1B_2$, then by intersecting the interior of π_6 . All points in the interior of the open segment G_1G_2 can see all interior points of π_n , but we cannot take a point on this segment as a Steiner point because of the coplanarity with some faces, however we are already close to it.

Now construct the Steiner point S as follows. Intersect the plane p_1 containing the arc $\widehat{A_1A_2}$ with the line passing through G_1 and G_2 . We obtain a point \tilde{S} in the interior of π_n . Now construct a plane p_2 which is orthogonal to the line passing through A_1 and A_2 and containing \tilde{S} . Intersect the planes p_1 and p_2 . We obtain a line which we will call l. By construction $\tilde{S} \in l$. Now take the point \tilde{S} and move it a little on the line l away from $\widehat{A_1A_2}$ but only so far, that it is still before the edge connecting B_1 with C_2 . The so obtained point is our Steiner point S, refer to Figure 4.



Fig. 4. The tetrahedron $A_1A_2G_1G_2$ in which the arc in Bagemihl's construction can lie is shown in blue. An example of such an arc $\widehat{A_1A_2}$ is also displayed in red. This arc cannot lie on a boundary face of the tetrahedron. The construction of a Steiner point S in π_n is also illustrated. The plane p_1 (not shown) contains the arc $\widehat{A_1A_2}$ and the plane p_2 (green) is orthogonal to the line passing A_1A_2 . One can see the line $l := p_1 \cap p_2$ and the points $P, \tilde{S}, S \in I$.

Proposition 2. A Bagemihl's polyhedron π_n with *n* vertices together with the constructed Steiner point $S \in \pi_n$ can be tetrahedralized.

Proof. By our construction, S lies beyond P. Recall that P is the intersection point of the tangent lines of the arc at A_1 and A_2 , see Figure 5. By that, the vertices $A_1, D_1, \ldots, D_k, A_2$ are visible by S (from the interior of π_n).

The visibility of the remaining vertices C_1, C_2, B_1 and B_2 is given, since S is chosen by moving \tilde{S} to the inside of the visible polytope of π_n . The point \tilde{S} lies on the boundary of the visible domain of the polytope π_n (and not only the polytope π_6). This polytope is as well bounded by the edge B_1C_2 , so we can be sure that there is space left in the interior.



Fig. 5. An example of the constructed Steiner point $S \in \pi_9$. The smaller polyhedron in the interior is the valid domain for π_9 .

2.3. Generalized Bagemihl's Polyhedra

Bagemihl gave the construction of a special class of polyhedra without leaving much space of freedom. The only choice one has is the one arc lying inside the original Schönhardt polyhedron which even has to fulfill some visibility constraints and the points one chooses on the arc. We can generalize these polyhedra without losing their main properties, i.e. they will still fulfill the properties (I)-(IV) of Theorem 1.

A similar way to generaize them is to let the rotation angle ϑ of the triangles be in $\vartheta \in (0^\circ, 60^\circ)$ instead of fixing it to a value of $\vartheta = 30^\circ$. At an angle of $\vartheta = 0^\circ$ the polyhedron is a prism and so doesn't fulfill the properties of the Theorem as well as in the case $\vartheta = 60^\circ$ in which the polyhedron separates into two parts which are attached at a single point, so the polyhedron isn't simple any more. Another way is to let the height *h* of the polyhedron be arbitrary in $h \in \mathbb{R}_{>0}$ instead of fixing it to the value of h = 1.

One can even change the bottom and top triangle itself. It is not necessary that they are parallel, are equilateral or have the same size. One can construct a generalized form of Bagemihl polyhedra based on two triangles in space connecting the vertices like in the original case. As long as they fulfill the properties of his Theorem and the open circular arc has the same visibility properties as in the original case, we will call them *generalized Bagemihl's polyhedra*. See Figure 6 for an example.

These polyhedra are still not decomposable, so at least one Steiner point is needed. But the construction of a Steiner point given above can be adapted to this class of polyhedra, so we can state the following Corollary. In Figure 6 the visible polytope of the example polyhedron is shown as well.

Corollary 3. For a tetrahedralization of $\tilde{\pi}_n$ with $n \in \mathbb{N}_{\geq 6}$ one needs exactly one Steiner point, where $\tilde{\pi}_n$ is a generalized Bagemihl's polyhedron as described above.

Proof. Since the property (II) mentioned in Theorem 1 is still fulfilled, at least one Steiner point is needed. On the other hand, one can use the construction as described in section 2.2 to obtain a Steiner point *S* in $\tilde{\pi}_n$. Furthermore, the proof of Proposition 2 can be adapted to show that *S* is already sufficient to tetrahedralize $\tilde{\pi}_n$.

3. A Larger Class of Indecomposable Polyhedra and The Number of Steiner Points

In this section, we first extend Bagemihl's Theorem to show that there exists a larger class of indecomposable polyhedra. We then provide a different construction of one of such polyhedra. We show that our constructed polyhedra are combinatorially the same as the Schönhardt and Bagemihl's polyhedra. But they may require more than one Steiner point to be decomposed. We then prove the maximum number of necessary Steiner points is $\left\lceil \frac{n-5}{2} \right\rceil$. Our proof is based on a construction of such Steiner points in the interior.



Fig. 6. Two views of an example of a generalized Bagemihl's polyhedron $\tilde{\pi}_6$. One still can add more vertices on the choosen suitable arc that is shown in red. The visible polytope is illustrated in a light blue, while the tetrahedron as the possible domain for arcs is illustrated in a darker blue to the left.

3.1. A Larger Class of Indecomposable Polyhedra

Note that Bagemihl's polyhedra π_n all satisfy a crucial property, which is (III) in Theorem 1, i.e., *every open* segment whose endpoints are vertices of π_n , but which is not an edge of σ_n , lies wholly exterior to π_n . This property is sufficient but not necessary to guarantee that a polyhedron is indecomposable. By relaxing this property, we can obtain a larger class of indecomposable polyhedra. Their properties are given in the following Theorem. The only difference to Theorem 1 is the property (III), which is also highlighted.

Theorem 4. If *n* is an integer not less than 6, then there exists a polyhedron, σ_n , with *n* vertices and the following properties:

- (I) σ_n is simple and every one of its faces is a triangle.
- (II) If τ is a tetrahedron, each of whose vertices is a vertex of σ_n , then not every interior point of τ is an interior point of σ_n .
- (III) Every open segment *e*, whose endpoints are vertices of σ_n , but which is not an edge of σ_n , satisfies $e \cap \sigma_n \neq e$.
- (IV) Every triangle whose sides are edges of σ_n is a face of σ_n .

Comment. The property (II) is redundant, since it can be derived from the other three properties. We keep it in order to keep the same form of Theorem 1.

The arguments that σ_n is an indecomposable polyhedron are exactly the same as those given by Bagemihl (see Section 2). The key fact is that every tetrahedron τ whose vertices in σ_n must contain at least one open segment. By the relaxed property (III), it is sufficient to ensure that some points of τ do not lie in the interior of σ_n .

3.2. A Construction of σ_n

Choose four non-coplanar points $\mathbf{a}, \mathbf{b}, \mathbf{c}, \mathbf{d} \in \mathbb{R}^3$, and a (simple) curve γ starting at \mathbf{c} and ending at \mathbf{d} , and γ lies in the intersection of the two open halfspaces bounded by the triangles **cda** and **dcb** (using the righthand-rule to oriented the triangles), refer to Figure 7 (a).

Now we will choose k + 2 ($k \ge 0$) distinct points, denoted as $\mathbf{g}_0, \ldots, \mathbf{g}_{k+1}$, on the curve γ from **c** to **d**, so that they all satisfy the following constraints (refer to Figure 7):

- (c1) The line segment **cd** intersects all the triangles \mathbf{abg}_i , $i = 0, \dots, k + 1$.
- (c2) Given two adjacent points \mathbf{g}_i and \mathbf{g}_{i+1} , for i = 0, ..., k, on the curve γ , the point \mathbf{g}_{i+1} and \mathbf{d} must lie in the same halfspace bounded by the plane containing \mathbf{abg}_i .

Author name / Procedia Engineering 00 (2015) 000-000

(1))	(a , c , d), (b , c , d)
(2))	$(a, c, g_0), (b, c, g_0), (a, d, g_{k+1}), (b, d, g_{k+1})$
(3))	$(a, g_i, g_{i+1}), (b, g_i, g_{i+1})$ where $i = 0,, k$

Table 1. The set of boundary faces of σ_n .

	a	b	c	d	\mathbf{g}_0	\mathbf{g}_1	g ₂	g ₃	g 4
x	-5	5	0	0	1	-2.5	2	-2.5	1
у	0	0	-10	10	-8	-4	0	4	8
z	0	0	2	2	4	4	4	4	4

Table 2. The coordinates of the vertices of a σ_9 .

- (c3) Let \mathbf{g}_i and \mathbf{g}_j , for i, j = -1, ..., k + 2 and $i \neq j$, be two non-adjacent points on the curve γ where $\mathbf{g}_{-1} := \mathbf{c}$ and $\mathbf{g}_{k+2} := \mathbf{d}$. Without loss of generality, assume i < j. Then the line segment $\mathbf{g}_i \mathbf{g}_j$ (except $\mathbf{g}_{-1} \mathbf{g}_{k+2} = \mathbf{cd}$) does not intersect all triangles \mathbf{abg}_l , where i < l < j.
- (c4) Let \mathbf{g}_i , \mathbf{g}_{i+1} and \mathbf{g}_{i+2} , for $i = -1, \dots, k$, be three consecutive points on the curve γ . Then the three points are neither coplanar with **a** nor **b**.

Now the polyhedron σ_n , n = 6 + k, where $k \ge 0$, is constructed by choosing the boundary faces listed in Table 1 (refer to Figure 7):

Comments. The curve γ in our construction is only an assistant. One can construct σ_n by simply choosing k points that satisfy all constraints. However, it is easier to imagine the relations of the points \mathbf{g}_i with a curve in mind. The condition (c4) ensures that all faces of σ_n are triangles.

Figure 7 gives an example of such polyhedron, σ_9 . A particular choice of the coordinates of the 9 vertices are given in the Table 2.



Fig. 7. An example polyhedron, σ_9 , with 9 vertices. The four initial points **a**, **b**, **c**, **d** as well as the points **g**_{*i*} *i* = 0,..., 4 chosen from a curve γ are shown in (a). Pictures (b), (c), and (d) are different views of the constructed polyhedron σ_9 . A particular choice of the coordinates of the vertices is given in Table 2.

In the following we show that the so constructed polyhedron σ_n satisfies the propertiers (I) - (IV) in Theorem 4. At first, we show that σ_n is a simple 3d polyhedron. Let \mathcal{T} be the set of tetrahedra

$$\mathcal{T} = \{\mathbf{abcg}_0, \mathbf{abdg}_{k+1}\} \cup \{\mathbf{abg}_i \mathbf{g}_{i+1} \mid i = 0, \dots, k\}.$$

The constraint (c2) ensures that every two tetrahedra in \mathcal{T} must either share a common face or only share the common edge **ab**. We see that the union of the set of tetrahedra of \mathcal{T} is a 3d simple polyhedron $P := \cup \mathcal{T}$. By constraint (c1), we see that the open line segment **cd** lies wholly in the interior of P. Moreover, this constraint also ensures that the two open triangles **cda** and **dcb** lie wholly in the interior of P. Finally, by removing the tetrahedron **abcd** from P we obtain the polyhedron σ_n .

Next we show that it is combinatorially equivalent to the Bagemihl's polyhedron π_n . The simplest case is when n = 6 (k = 0). The corresponding π_6 is just the well-known Schönhardt polyhedron (refer to Figure 1). The 6 vertices of σ_6 are: **a**, **b**, **c**, **d**, **g**₀, **g**₁, respectively. We map them ono-to-one to the vertices of the Schönhardt polyhedron as following (see Figure 8 Left):

•
$$\mathbf{g}_0 \to A_1, \mathbf{c} \to B_1, \mathbf{b} \to C_1$$
; and

•
$$\mathbf{g}_1 \to A_2, \mathbf{a} \to B_2, \mathbf{d} \to C_2.$$

In general, when $n \ge 6$ ($k \ge 0$), the *n* vertices of σ_n are: **a**, **b**, **c**, **d**, **g**₀, **g**₁, ..., **g**_{k+1}, respectively. We build a one-to-one map between the vertices of σ_n and the Bagemihl's polyhedron π_n as following (refer to Figure 8 Right):

- $\mathbf{g}_0 \to A_1, \mathbf{c} \to B_1, \mathbf{b} \to C_1;$
- $\mathbf{g}_{k+1} \rightarrow A_2, \mathbf{a} \rightarrow B_2, \mathbf{d} \rightarrow C_2$; and
- $\mathbf{g}_i \to D_i$, for all $i = 1, \ldots, k$.

By this mapping, one can check that the faces of σ_n and π_n are also mapped one-to-one, so do the edges of them.



Fig. 8. The mapping between the vertices of σ_n and the vertices of the Bagemihl's polyhedron π_n .

Now we can show that σ_n satisfies the properties in Theorem 4 by borrowing Bagemihl's arguments in proving Theorem 1 [1].

At first, σ_n satisfies the properties (I) and (IV) by a direct checking of the faces (listed above) and the edges of σ_n , using (c3) and (c4). Note that the property (IV) is fulfilled because of the constraint (c3) which prohibits to have three consecutive points $\mathbf{g}_i, \mathbf{g}_{i+1}, \mathbf{g}_{i+2}$ being collinear and constraint (c4) ensures the triangularity of the faces.

Remember that the main reason that causes σ_n to be indecomposable is the property (II), since no open tetrahedron τ whose vertices of σ_n can lie in the interior of σ_n . The key fact is that τ must contain an open segment as stated in the property (III). This fact is true by the properties (I), (III) and (IV).

What remains is to show that σ_n satisfies the property (III). The open segments of σ_n are the line segment **ab**, and all line segments with endpoints $\mathbf{g}_i \mathbf{g}_j$, where \mathbf{g}_i and \mathbf{g}_j are not adjacent vertices on the curve γ , where $i, j = -1, 0, \dots, k+2$ and $i \neq j$, except the line segment $\mathbf{cd} = \mathbf{g}_{-1}\mathbf{g}_{k+2}$. The constraint (c3) in our construction ensures that such a line segment must not lie wholly in the interior of σ_n . This shows that the property (III) is satisfied.

3.3. The Number of Steiner Points for σ_n

From now on, we study the question: "Given a σ_n , how many Steiner points are necessary to decompose it?"

At first, we show that a σ_n may need more than one Steiner point to be decomposed. We provide an explicit example of a σ_9 . The geometry of this polyhedron is similar to the one shown in Figure 7. The coordinates of the 9 vertices are given in the Table 2.

Given an arbitrary σ_n , we can associate every pair of its triangles, $\mathbf{g}_i \mathbf{g}_{i+1} \mathbf{a}$ and $\mathbf{g}_i \mathbf{g}_{i+1} \mathbf{b}$, to an *interval*, $\mathbf{t}_{a,i} \mathbf{t}_{b,i}$, on the line through **cd**, for all i = 0, 1, ..., k, where:

$$\mathbf{t}_{a,i} := \text{plane}_{\mathbf{g}_i \mathbf{g}_{i+1} \mathbf{a}} \cap \text{line}_{\mathbf{cd}}$$
$$\mathbf{t}_{b,i} := \text{plane}_{\mathbf{g}_i \mathbf{g}_{i+1} \mathbf{b}} \cap \text{line}_{\mathbf{cd}}.$$

Any point **p** in the interval $\mathbf{t}_{a,i}\mathbf{t}_{b,i}$ must see the two triangles, $\mathbf{g}_i\mathbf{g}_{i+1}\mathbf{a}$ and $\mathbf{g}_i\mathbf{g}_{i+1}\mathbf{b}$, from the interior of σ_n . However, if **p** does not lie in the interval defined by a pair of triangles, **p** can not see them.

In general, such an interval is not necessarily inside the line segment **cd** for an arbitrary σ_n . In our particular example (in Table 2), the pair of planes containing the triangles $\mathbf{g}_i \mathbf{g}_{i+1} \mathbf{a}$ and $\mathbf{g}_i \mathbf{g}_{i+1} \mathbf{b}$ cut the line segment **cd** in its interior, for all i = 0, 1, 2, 3. Furthermore, the two intervals $\mathbf{t}_{a,0} \mathbf{t}_{b,0}$ and $\mathbf{t}_{a,3} \mathbf{t}_{b,3}$ are disjoint. This implies that one cannot find a common interior Steiner point that is visible simultaneously by the four triangles, which are $\mathbf{g}_0 \mathbf{g}_1 \mathbf{a}$, $\mathbf{g}_0 \mathbf{g}_1 \mathbf{b}, \mathbf{g}_3 \mathbf{g}_4 \mathbf{a}$, and $\mathbf{g}_3 \mathbf{g}_4 \mathbf{b}$. Hence, more than one Steiner point is needed for decomposing this polyhedron.

For a general σ_n with *n* vertices, there are k + 1 intervals. It is easy to estimate that the required number of Steiner points for decomposing σ_n will not exceed the total number of such intervals, which is k + 1 = n - 5. However, this estimate is too rough. By an careful construction of Steiner points, one can get an explicit upper bound on the number of Steiner points for any σ_n . It is given by the following Theorem.

Theorem 5. Given a 3d polyhedron σ_n with n vertices, it can be decomposed by adding $\left[\frac{n-5}{2}\right]$ interior Steiner points.

Proof. We prove it in two steps: at first, we will place this number of Steiner points in the interior of the σ_n , then we show how to tetrahedralise it with these Steiner points.

Step (1), placing Steiner points. From the previous analysis, we see that the requirement of multiple Steiner points comes from the fact that one may not find a common Steiner point that is simultaneously visible by all boundary faces. We will place a number of Steiner points in the interior of σ_n . We make sure that each Steiner point that we place will be visible by a certain number of boundary faces, and every boundary face will be visible by at least one of these Steiner points.

Consider the k + 2 vertices, $\mathbf{g}_0, \mathbf{g}_1, \dots, \mathbf{g}_{k+1}$ of σ_n . Each \mathbf{g}_i ($i = 0, 1, \dots, k+1$) is a vertex of four adjacent boundary faces of σ_n , i.e., $\mathbf{g}_{i-1}\mathbf{g}_i\mathbf{a}, \mathbf{g}_{i-1}\mathbf{g}_i\mathbf{b}, \mathbf{g}_i\mathbf{g}_{i+1}\mathbf{a}$, and $\mathbf{g}_i\mathbf{g}_{i+1}\mathbf{b}$ (recall that $\mathbf{g}_{-1} = \mathbf{c}$ and $\mathbf{g}_{k+2} = \mathbf{d}$). We will search a point inside σ_n and near to \mathbf{g}_i , hence it is visible by all these four faces. For this purpose, it is not necessary to use all \mathbf{g}_i . In particular, we choose the following subset of the set of vertices of σ_n ,

$$\mathcal{G} := \{\mathbf{g}_1, \mathbf{g}_3, \mathbf{g}_5, \ldots, \mathbf{g}_m\},\$$

where the last index is

$$m = k + ((k + 1) \text{ modulo } 2),$$

which is the largest odd number of the indices. The cardinality $|\mathcal{G}| = \left[\frac{k+1}{2}\right]$.

Let \mathcal{G}^* be the set of points constructed from the points in \mathcal{G} , such that

$$\mathcal{G}^* := \{\mathbf{g}_i^* := \text{plane}_{\mathbf{g}, \mathbf{ab}} \cap \text{line}_{\mathbf{cd}} \mid \mathbf{g}_i \in \mathcal{G}\},\$$

see Figure 9. Note that the points in \mathcal{G}^* have the following visibilities from the inside of σ_n :

- (i) Each \mathbf{g}_{i}^{*} , where j = 1, 3, ..., m, is visible by the faces $\mathbf{ag}_{j-1}\mathbf{g}_{j}$, $\mathbf{bg}_{j-1}\mathbf{g}_{j}$, $\mathbf{ag}_{j}\mathbf{g}_{j+1}$, and $\mathbf{bg}_{j}\mathbf{g}_{j+1}$;
- (ii) Additionally, \mathbf{g}_1^* is visible by the two faces \mathbf{acg}_0 and \mathbf{bcg}_0 , and \mathbf{g}_m^* , is visible by the two faces \mathbf{adg}_{k+1} and \mathbf{bdg}_{k+1} .



Fig. 9. Two views of the example polyhedron σ_9 with the constructed points \mathcal{G}^* and the Steiner points \mathcal{S} on the common line *l* that is parallel to cd.

(1)	(a , c , d), (b , c , d);
(2)	$(\mathbf{a},\mathbf{c},\mathbf{s}_1),(\mathbf{b},\mathbf{c},\mathbf{s}_1),(\mathbf{a},\mathbf{d},\mathbf{s}_m),(\mathbf{b},\mathbf{d},\mathbf{s}_m);$
(3)	{ $(\mathbf{a}, \mathbf{s}_p, \mathbf{s}_{p+2}), (\mathbf{b}, \mathbf{s}_p, \mathbf{s}_{p+2}) \mid p = 1, 3, 5, \dots, m$ }

Table 3. The set of boundary faces of the remaining polyhedron.

Since all \mathbf{g}_i^* 's are on the line segment \mathbf{cd} , i.e., they are coplanar with the two faces \mathbf{cda} and \mathbf{cdb} , they are not yet our wanted Steiner points.

Now we will place our Steiner points moving from the set of points \mathcal{G}^* . Choose the plane p along the middle axis of the two planes containing the two bottom triangles cda and cdb. p must contain the line segment cd. Now we will move \mathcal{G}^* within the plane p and toward the interior of σ_n . For our proof, we just move all points in such a way, that they all stay within a line, denoted as l, which is parallel to **cd**. And we choose the moving distance small enough such that all the moving points remain in the interior of σ_n and their visibilities by faces given in (i) and (ii) do not change. We then take the set of points on the line as our Steiner points, denoted as

$$S := \{\mathbf{s}_i \mid \mathbf{s}_i \text{ is moved from } \mathbf{g}_i^* \text{ onto } l, \ \mathbf{g}_i^* \in \mathcal{G}^*\},\$$

then $|S| = \left\lceil \frac{k+1}{2} \right\rceil$. See Figure 9 for an example. Step (2), tetrahedralising σ_n . With the created Steiner points, we are able to create a tetrahedralisation of σ_n . The idea is first to remove tetrahedra from σ_n by using the set of Steiner points and the visibility properties of them, and second to tetrahedralise the remaining part, which is a 3d polyhedron with vertices **a**, **b**, **c**, **d**, and the set of Steiner points.

By the visibility properties (i) and (ii), we can remove the following sets of tetrahedra from σ_n :

$$\mathcal{T}_1 := \{ \mathbf{ag}_{j-1} \mathbf{g}_j \mathbf{s}_j, \mathbf{bg}_{j-1} \mathbf{g}_j \mathbf{s}_j, \mathbf{ag}_j \mathbf{g}_{j+1} \mathbf{s}_j, \mathbf{bg}_j \mathbf{g}_{j+1} \mathbf{s}_j | \mathbf{s}_j \in \mathcal{S} \}, \\ \mathcal{T}_2 := \{ \mathbf{acg}_0 \mathbf{s}_1, \mathbf{bcg}_0 \mathbf{s}_1, \mathbf{adg}_{k+1} \mathbf{s}_m, \mathbf{bdg}_{k+1} \mathbf{s}_m \}, \\ \mathcal{T}_3 := \{ \mathbf{ag}_{p+1} \mathbf{s}_p \mathbf{s}_{p+2}, \mathbf{bg}_{p+1} \mathbf{s}_p \mathbf{s}_{p+2} | p = 1, 3, 5, \dots, m-2 \}.$$



Fig. 10. The polyhedron after removing the sets $\mathcal{T}_1, \mathcal{T}_2$ and \mathcal{T}_3 of tetrahedra from σ_n . A decomposition of this polyhedron into the two sets \mathcal{T}_4 and \mathcal{T}_5 of tetrahedra is given. The internal edges are shown in brown.

The remaining region of σ_n after removing the above tetrahedra is a 3d polyhedron, see Figure 10 for an example. It has the vertices $\{a, b, c, d\} \cup S$, and the boundary faces are given in Table 3. Since all Steiner points are collinear, this polyhedron can be decomposed into the following two sets of tetrahedra:

$$\mathcal{T}_4 := \{ \mathbf{acds}_1, \mathbf{bcds}_1 \},$$

$$\mathcal{T}_5 := \{ \mathbf{as}_p \mathbf{s}_{p+2} \mathbf{d}, \mathbf{bs}_p \mathbf{s}_{p+2} \mathbf{d} \mid p = 1, 3, 5, \dots, m-2 \}$$

This concludes our proof.

There are indeed many possibilities to place Steiner points that may lead to a smaller number of Steiner points. However, we can show that this number $\left\lceil \frac{k+1}{2} \right\rceil = \left\lceil \frac{n-5}{2} \right\rceil$ of Steiner points is optimal in the worst case.

Theorem 6. Given $n \in \mathbb{N}_{\geq 6}$, one can construct an a 3d polyhedron σ_n with n vertices which has the property that one needs exactly $\left\lfloor \frac{n-5}{2} \right\rfloor$ interior Steiner points to decompose it.

Proof. We prove the Theorem by giving a general construction of a σ_n , so that it will always need at least this number of Steiner points. We then get the equality by Theorem 5. The basic idea is to control the overlap of the intervals $\mathbf{t}_{a,j}\mathbf{t}_{b,j}$, j = 0, ..., n - 6, as defined at the beginning of section 3.3. By the following construction we ensure that two consecutive intervals $\mathbf{t}_{a,j}\mathbf{t}_{b,j+1}$ and $\mathbf{t}_{a,j+1}\mathbf{t}_{b,j+1}$ overlap in their interior. Moreover, if two non-consecutive intervals don't overlap, we will obtain the desired number of Steiner points.

Fix $n \ge 6$ and start with non-coplanar points $\mathbf{a}, \mathbf{b}, \mathbf{c}, \mathbf{d} \in \mathbb{R}^3$ as described in section 3.2. Then choose n - 4 points \mathbf{g}_i , i = 0, ..., n - 5 from the valid domain of a curve in a zig-zag shape, like in the polyhedron in Figure 7. By moving the points \mathbf{g}_i lower, so that the segments $\mathbf{g}_i \mathbf{a}$ or $\mathbf{g}_i \mathbf{b}$ resp. are nearly crossing the line \mathbf{cd} , we obtain non overlapping intervals $\mathbf{t}_{a,i}\mathbf{t}_{b,i}$ and $\mathbf{t}_{a,i+2}\mathbf{t}_{b,i+2}$. So, we can achieve that except for the consecutive intervals $\mathbf{t}_{a,j}\mathbf{t}_{b,j}$ and $\mathbf{t}_{a,j+1}\mathbf{t}_{b,j+1}$ with j = 0, ..., n - 6, no intervals overlap. Placing one Steiner point slightly above each overlap of the intervals gives the number of $\left\lfloor \frac{n-5}{2} \right\rfloor$ interior Steiner points. One can decompose the polyhedron as described in the proof of Theorem 5. \Box



Fig. 11. An example polyhedron, σ_{12} , with 12 vertices. The coordinates of the vertices are given in the Table 4. Different views of this polyhedron are shown in (a), (b), (c), and (d), respectively. In particular, two overlappings intervals are shown in (c).

Figure 11 shows a particular example of such a polyhedron with 12 vertices. The coordinates of the 12 vertices is given in Table 4. By our construction, this polyhedron satisfies the property that only two adjacent intervals are overlapping. A pair of such intervals is illustrated in Figure 11 (c). Therefore, this polyhedron needs at least 4 Steiner points to be decomposed, which is optimal for this case.

Author name / Procedia Engineering 00 (2015) 000-000

	a	b	c	d	g 0	\mathbf{g}_1	g ₂	g ₃
x	-1.294	4.830	4.830	-3.536	4.253	-0.301	3.117	-2.183
y	10	0	10	0	6.532	9.760	2.999	8.657
z	4.830	1.294	-1.294	3.536	-2.426	0	-2.571	0.646
	g 4	g 5	g 6	g 7				
x	1.874	-3.330	0.163	-4.051				
у	1.002	6.864	-0.105	3.184				
z	-1.808	1.350	-0.366	2.242				

Table 4. A choice of the coordinates of the vertices of a σ_{12} . The geometry of this polyhedron is shown in Figure 11. With these coordinates, this polyhedron needs at least 4 Steiner points to be decomposed.

4. Discussion

We comment that our construction of a Steiner point in Bagemihl's polyhedra as well as it generalizations is not unique. Note that the line segment B_1C_2 in the Bagemihl's polyhedron always touches the visible polyhedron. It is always possible to choose a Steiner point near this line segment such that it can decompose the polyhedron. The Steiner point can be chosen such that it is near the interval cut by the two planes containing the triangles $A_1A_2C_1$ and $A_1A_2B_2$ for π_6 (or $A_1A_2D_1$ and $D_kA_2B_2$ for π_n , where n > 6) and the line segment B_1C_2 .

The Theorem 1 of Bagemihl proves the existence of a class of polyhedra (that satisfy all these properties). The following two open questions are interesting:

- (1) Except the generalized Bagemihl's polyhedra $\tilde{\pi}_n$, is there other construction that satisfies Theorem 1.
- (2) Does every polyhedron satisfying the properties in Theorem 1 require only one Steiner point to be decomposed?

The result of Theorem 5 shows that any σ_7 needs only one Steiner points to be decomposed, regardless of its geometry. However, we do not know whether this is true or false for an arbitrary 3d polyhedron with 7 vertices.

Acknowledgments

This work is supported under the research project "TetGen" in Weierstrass Institute for Applied Analysis and Stochastics (WIAS), Berlin. The authors sincerely thank the reviewers for their precious comments and suggestions.

References

- [1] Bagemihl, F. (1948). On indecomposable polyhedra. The American Mathematical Monthly, 55(7):411-413.
- [2] Below, A. (2002). Complexity of Triangulation. PhD thesis, Swiss Federal Institute of Technology, ETH Zürich. DISS. ETH No. 14672, 2002.
- [3] Chazelle, B. (1984). Convex partition of polyhedra: a lower bound and worst-case optimal algorithm. SIAM Journal on Computing, 13(3):488– 507.
- [4] De Loera, J. A., Rambau, J., and Santos, F. (2010). *Triangulations, Structures for Algorithms and Applications*, volume 25 of *Algorithms and Computation in Mathematics*. Springer Verlag Berlin Heidelburg.
- [5] George, P., Borouchaki, H., and Saltel, E. (2003). Ultimate robustness in meshing an arbitrary polyhedron. *International Journal for Numerical Methods in Engineering*, 58:1061–1089.
- [6] George, P., Hecht, F., and Saltel, E. (1991). Automatic mesh generator with specified boundary. *Computer Methods in Applied Mechanics and Engineering*, 92:269–288.
- [7] Lennes, N. (1911). Theorems on the simple finite polygon and polyhedron. American Journal of Mathematics, 33(1/4):37–62.
- [8] Rambau, J. (2005). On a generalization of Schönhardt's polyhedron. In Goodman, J. E., Pach, J., and Welzl, E., editors, *Combinatorial and Computational Geometry*, volume 52, pages 501–516. MSRI publications.
- [9] Ruppert, J. and Seidel, R. (1992). On the difficulty of triangulating three-dimensional nonconvex polyhedra. Discrete & Computational Geometry, 7:227–253.
- [10] Schönhardt, E. (1928). Über die Zerlegung von Dreieckspolyedern in Tetraeder. Mathematische Annalen, 98:309-312.
- [11] Si, H. (2015). TetGen, a Delaunay-based quality tetrahedral mesh generator. ACM Trans. Math. Softw., 41(2):11:1–11:36.
- [12] Weatherill, N. P. and Hassan, O. (1994). Efficient three-dimensional Delaunay triangulation with automatic point creation and imposed boundary constraints. *International Journal for Numerical Methods in Engineering*, 37:2005–2039.