


Global Sensitivity Analysis Using the Ultra-Low Resolution Energy Exascale Earth System Model (E3SM)




Irina Tezaur, Kara Peterson, Amy Powell, John Jakeman, Erika Roesler
Sandia National Laboratories, Livermore, CA and Albuquerque, NM

European Seminar on COmputing (ESCO) 2022
Pilsen, CA June 13-17, 2022

SAND2022-3734C

- Background and motivation
- Methodology
- Model tuning and spin-up
- Evaluation of the ultra-low resolution (ULR) E3SM
- Global sensitivity analysis results
- Summary




The focus of this talk is the following paper:

- Paper is **under review** for publication in the *Journal of Advances in Modeling Earth Systems (JAMES)*.
- Pre-print **available** at <https://www.essoar.org/doi/10.1002/essoar.10508267.2>

The screenshot shows the ESSOAr (Earth and Space Science Open Archive) website. At the top, there is a navigation bar with links for 'ABOUT US', 'SUBMIT', and 'AUTHOR DASHBOARD'. Below this, a banner indicates that the preprint has been submitted to and is under consideration at the Journal of Advances in Modeling Earth Systems (JAMES). The main title of the preprint is 'Global sensitivity analysis using the ultra-low resolution Energy Exascale Earth System Model'. The authors listed are Irina Kalashnikova Tezaur, Kara Peterson, Amy Powell, John Jakeman, and Erika Roesler. The publication date is Thursday, 14 Oct 2021, with a DOI link provided. Below the title and authors, there are buttons for 'Download PDF' and 'Cite', along with 'Tools' and 'Share' options. The abstract section begins with the text: 'For decades, the Arctic has been warming at least twice as fast as the rest of the globe. As a first step towards quantifying parametric uncertainty in Arctic feedbacks, we perform a variance-based global sensitivity analysis (GSA) using a fully-coupled, ultra-low resolution (ULR) configuration of version 1 of the Department of Energy's Energy Exascale Earth System Model (E3SMv1). The study randomly draws 139 realizations of ten model parameters spanning three E3SMv1 components (sea ice, atmosphere and ocean), which are used to generate 75 year long projections of future climate using a fixed pre-industrial forcing. We quantify the sensitivity of six Arctic-focused quantities of interest (QOIs) to these parameters using main effect, total effect and Sobol sensitivity indices computed with a Gaussian process emulator. A sensitivity index-based ranking of model parameters shows that the atmospheric parameters in the CLUBB (Cloud Layers Unified by Binormals) scheme have significant impact on sea ice status and the larger Arctic climate. We also use the Gaussian process emulator to predict the response of varying each variable when the impact of other parameters are averaged out. These results allow one to assess the non-linearity of a parameter's impact on a QOI and investigate the presence of local minima encountered during the spin-up tuning process. Our study confirms the necessity of performing global analyses involving fully-coupled climate models, and motivates follow-on investigations in which the ULR model is compared rigorously to higher resolution configurations to confirm its viability as a lower-cost surrogate in fully-coupled climate uncertainty analyses.'






- **Background and motivation**
- Methodology
- Model tuning and spin-up
- Evaluation of the ultra-low resolution (ULR) E3SM
- Global sensitivity analysis results
- Summary



Background and motivation

Arctic systems are **strongly coupled and rapidly changing!**

- Arctic is **warming** at twice the rate of the rest of the globe¹
- Abrupt changes related to **sea ice loss**, **land ice melt** and **permafrost thaw** have the potential to cause significant **global climate impacts**²
 - September **sea ice extent** has declined **13.1% per decade** from 1979-2020 relative to 1979-2010 average¹
 - **Global mean sea-level** is rising at the rate of **3.2 mm/year** and this rate is increasing due to **melting of polar ice sheets**
 - **Arctic permafrost** erosion rates have **accelerated**, leading to threats to coastal communities, coastal infrastructure and global carbon balance
- Changes can potentially lead to **tipping events** with significant global impacts²
- Research to **advance predictability** and **bound uncertainty** of Earth system models are **crucial** to inform **planning** and **decision-making**



Gibbs & Richmond, 2015

¹ NOAA Arctic Report Card 2020.

² "Climate tipping points - too risky to bet against", Lenton *et al.* Nature 2019.

6 Objective and Approach

Objective:

- Gain understanding of **Arctic system dynamics** including feedbacks between Arctic physical systems using **sea ice** as an **exemplar**


Approach:

- Perform **global sensitivity analysis (GSA)** using the **fully-coupled ultra-low resolution (ULR)** configuration of the Energy Exascale Earth System Model (E3SM)
 - Investigate **uncertainty** and **stability** in E3SM simulations
 - **Analyze trends** and investigate **internal variability** in E3SM simulations of sea ice extent


Energy Exascale Earth System Model (E3SM):

- U.S. DOE flagship **open-source**¹ coupled Earth system model
- **Collaboration** between 8 national labs and 12 academic institutions
- Development driven by **DOE Office of Science mission interests**: energy and water issues looking out 40 years
- Our study utilized version 1 of the E3SM, denoted by **E3SMv1**, with **pre-industrial control (1850)** forcing at the **ULR** configuration

¹ www.e3sm.org; <https://github.com/E3SM-Project/E3SM>.



- Background and motivation
- **Methodology**
- Model tuning and spin-up
- Evaluation of the ultra-low resolution (ULR) E3SM
- Global sensitivity analysis results
- Summary



Global Sensitivity Analysis (GSA): find which parameters have largest impact on model's QOIs

- Initial step in **quantifying uncertainty** in model parameterizations and measuring their impact on QOIs
- **Pros:** considers parameter sensitivity over **entire domain**, IDs **cross-component parameter interactions**
- **Cons:** computationally **expensive!**
- **Sobol sensitivity analysis:** quantifies global sensitivity of a QOI (f) as the fraction of the variance due to each parameter (z_i)

ANOVA Expansion of QOI:
$$f(z) = \hat{f}_0 + \sum_{i=1}^d \hat{f}_i(z_i) + \sum_{i,j=1}^d \hat{f}_{i,j}(z_i, z_j) + \sum_{i,j,k=1}^d \hat{f}_{i,j,k}(z_i, z_j, z_k) + \dots = \sum_{u \subseteq D} \hat{f}_u(z_u)$$

Variance of QOI:
$$\mathbb{V}[f] = \sum_{u \subseteq D} \mathbb{V}[\hat{f}_u]$$

Main effect indices:
$$S_i^M = \frac{\mathbb{V}[\hat{f}_{e_i}]}{\mathbb{V}[f]}$$
 Measure effect of individual parameters acting alone

Total effect indices:
$$S_i^T = \frac{\sum_{u \subseteq J} \mathbb{V}[\hat{f}_u]}{\mathbb{V}[f]}$$

Sobol indices:
$$S_u = \frac{\mathbb{V}[\hat{f}_u]}{\mathbb{V}[f]}$$

Measure contribution of each parameter and parameter interactions to the variance of QOI



Component	Notation	Parameter	Units	Min	Max	Description
Sea Ice (MPAS-Sealce)	z_1	ksno	W/(mK)	0.2	0.6	Snow conductivity
	z_2	lambda_pond	1/s	1.15e-8	1.15e-4	Drainage timescale of ponds
	z_3	dragio	—	0.2e-3	160e-3	Neutral ocean-ice drag
Atmosphere ¹ (EAM)	z_4	clubb_c1	—	1.0	5.0	Const assoc. w/ dissipation of variance w'^2
	z_5	clubb_c8	—	2.0	8.0	Const assoc. w/ Newtonian damping of w'^3
	z_6	gamma_coeff	—	0.1	0.5	Const of width of PDF in w coord
	z_7	cldfrc_dp1	—	0.02	0.1	Deep convection cloud fraction parameter
Ocean (MPAS-O)	z_8	standardgm_tracer_kappa	m ² /s	600	1800	Bolus coefficient of GM parameterization of eddy transport
	z_9	cvmix_kpp_criticalbulkrichardson_number	—	0.2	1.0	Bulk Richardson number used in KPP vertical mixing scheme

¹ CLUBB (Cloud Layers Unified by Binormals): cloud physics parameterization in EAM.



- **Nine parameters span three E3SM components** (sea ice, atmosphere, ocean)
- Parameters and parameter ranges were guided by **past analyses** [Urrego-Blanco *et al.*, 2016; Urrego-Blanco *et al.* 2019; Reckinger *et al.*, 2015; Asay-Davis *et al.*, 2018; Qian *et al.*, 2018; Rasch *et al.*, 2019]




QOI	Units	Description	Component
SIE	km ²	Total Arctic sea ice extent	Sea ice
SIV	km ³	Total Arctic sea ice volume	Sea ice
SST	°C	Sea surface temperature averaged over 60-90° N	Ocean
TS	°C	Surface air temperature averaged over 60-90° N	Atmosphere
FLNS	W/m ²	Net longwave flux at surface over 60-90° N	Atmosphere
CLDLow	—	Low cloud coverage below 700 hPa averaged over 60-90° N	Atmosphere

- Six QOIs span three E3SM components (sea ice, ocean and atmosphere)
- QOIs are **Arctic-focused**, motivated by literature [Urrego-Blanco *et al.*, 2016; Urrego-Blanco *et al.*, 2019]
- QOIs are computed by **averaging annually** and **seasonally** over last 25 years of each 75-year simulation

GSA workflow with fully-coupled E3SM



Above: ULR atmosphere grid ($\approx 7.5^\circ$)
Below: ULR ocean/sea ice grid (240km or $\approx 2.2^\circ$)




The **ULR configuration** enables sufficient ensemble generation for **full GSA** with the **fully-coupled E3SM!**

- ULR E3SM configuration is $\approx 100 \times$ less expensive to run than the “standard” 1° resolution E3SM
 - Using the ULR E3SM, our study took $\approx 1.00 \times 10^6$ CPU hours¹ on Sandia’s Skybridge HPC cluster
 - The same study using the standard 1° resolution E3SM would require $\approx 1.14 \times 10^8$ CPU hours

¹ Equivalent to 24 sydp on 6 Skybridge nodes.

- Background and motivation
- Methodology
- **Model tuning and spin-up**
- Evaluation of the ultra-low resolution (ULR) E3SM
- Global sensitivity analysis results
- Summary



13 Model tuning and spin-up

Objective: before initializing perturbed runs, run the model until an **equilibrium, Earth-like state** is achieved using pre-industrial (1850) control (piControl) forcing in which we have


- Constant global average mean **surface air temperature**, (a)
- Near-zero long-term average net **top-of-atmosphere (TOA) energy flux**, (b)
- Stable yearly **sea ice coverage**, (c)

Spin-up approach:


- 500 year run with pre-industrial control forcing, **default parameter values**
- 180 year run branched from 500 year initial run, with **parameter values from Golaz *et al.* 2018**
- **Year 675** of the branch run was used as **IC** for GSA runs

Right figures: bold lines indicate linear trends in years 26-500 and 526-800. Trends are much closer to 0 for branch run (slope = -0.00082 , 0.0005 , 0.0012 for surface temperature, TOA flux, Arctic sea ice, respectively)


(a) Global Yearly Average Surface Temperature



(b) Global Yearly Average Net Flux at Atmosphere




(c) Global Yearly Average Sea Ice Extent






- Background and motivation
- Methodology
- Model tuning and spin-up
- **Evaluation of the ultra-low resolution (ULR) E3SM**
- Global sensitivity analysis results
- Summary



How good is the ULR E3SM?



- ULR configuration of the E3SM has not been **scientifically validated**; it was designed primarily for **rapid turn-around testing**
- Years 526-675 of the ULR simulation were compared to **observational data** from CERES-EBAF Ed4.1 [Loeb *et al.* 2018] and ERA-Interim reanalysis [Dee *et al.*, 2011], and **1° resolution E3SM** simulation data



Left figures: TOA flux (W/m^2) for (a) years 526-675 of branched ULR spin-up simulation and (b) years 1-500 of the 1° piControl, compared with CERES-EBAF Ed4.1 data

Although ULR simulation does not capture small-scale/regional features seen in 1° resolution simulation, large-scale patterns are similar.

How good is the ULR E3SM?




- ULR configuration of the E3SM has not been **scientifically validated**; it was designed primarily for **rapid turn-around testing**
- Years 526-675 of the ULR simulation were compared to **observational data** from CERES-EBAF Ed4.1 [Loeb *et al.* 2018] and ERA-Interim reanalysis [Dee *et al.*, 2011], and **1° resolution E3SM** simulation data

Model

ERA-Interim

Diff.



Left figures: Total precipitation (mm/day) for (a) years 526-675 of branched ULR spin-up simulation and (b) years 1-500 of the 1° piControl, compared with ERA-Interim data


Although ULR simulation does not capture small-scale/regional features seen in 1° resolution simulation, large-scale patterns are similar.

17 How good is the ULR E3SM?




- ULR configuration of the E3SM has not been **scientifically validated**; it was designed primarily for **rapid turn-around testing**
- Years 526-675 of the ULR simulation were compared to **observational data** from CERES-EBAF Ed4.1 [Loeb *et al.* 2018] and ERA-Interim reanalysis [Dee *et al.*, 2011], and **1° resolution E3SM** simulation data

Model




Left figures: Zonal temperature (°C) for (a) years 526-675 of branched ULR spin-up simulation and (b) years 1-500 of the 1° piControl, compared with ERA-Interim data, to demonstrate vertical variation in atmosphere

ERA-Interim



Diff.






Temperature field shows most divergence from observations (warm bias of ~8°C)

(a) ULR E3SM

(b) 1° E3SM

- Background and motivation
- Methodology
- Model tuning and spin-up
- Evaluation of the ultra-low resolution (ULR) E3SM
- **Global sensitivity analysis results**
- Summary





- Most perturbed runs have reached **equilibrium** by **year 40**
- Runs exhibit a great deal of **variability**:
 - Several runs result in complete loss of Arctic sea ice
 - Several runs exhibit apparent exponential growth in sea ice

Figures above: “Spaghetti plot” of two QOIs for our 139 successful ULR runs. Red markers indicate baseline run (no parameter perturbation)

GSA results: sensitivity indices


¹ Measure effect of individual parameters acting alone

² Measure variance that remains after learning values of every variable except z_i


³ Measure contribution of interaction between parameter subset u on the variance of a QOI




Main Effects¹



Sobol Indices²



Total Effects³



Sea ice
params

Atmosphere
params

Ocean
params

- Figure above shows sensitivity indices for Sea Ice Volume QOI in spring

GSA results: sensitivity indices


¹ Measure effect of individual parameters acting alone

² Measure variance that remains after learning values of every variable except z_i

³ Measure contribution of interaction between parameter subset u on the variance of a QOI



Main Effects¹




Sea ice
params


Atmosphere
params

Ocean
params

Sobol Indices²



Total Effects³



- Figure above shows sensitivity indices for Sea Ice Volume QOI in spring
- Atmospheric parameters related to cloud parameterizations (CLUBB) are most important for all QOIs; z_6 (clubb_c8) has largest total effects

GSA results: sensitivity indices


¹ Measure effect of individual parameters acting alone

² Measure variance that remains after learning values of every variable except z_i


³ Measure contribution of interaction between parameter subset u on the variance of a QOI




Main Effects¹



Sobol Indices²



Total Effects³



Sea ice
params

Atmosphere
params

Ocean
params

z_5 (clubb_c1) and z_6 (clubb_c8) combined effects are stronger than total effects for 5 insensitive parameters

- Figure above shows sensitivity indices for Sea Ice Volume QOI in spring
- Atmospheric parameters related to cloud parameterizations (CLUBB) are most important for all QOIs; z_6 (clubb_c8) has largest total effects
- Analysis also reveals significant parameter interactions between atmosphere parameters

GSA results: sensitivity indices


¹ Measure effect of individual parameters acting alone

² Measure variance that remains after learning values of every variable except z_i


³ Measure contribution of interaction between parameter subset u on the variance of a QOI




Main Effects¹



Sobol Indices²



Total Effects³



Sea ice
params

Atmosphere
params

Ocean
params

z_5 (clubb_c1) and z_6 (clubb_c8) combined effects are stronger than total effects for 5 insensitive parameters

- Figure above shows **sensitivity indices** for **Sea Ice Volume QOI** in **spring**
- Atmospheric parameters related to cloud parameterizations (CLUBB) are **most important** for all QOIs; z_6 (clubb_c8) has largest total effects
- Analysis also reveals **significant parameter interactions** between atmosphere parameters
- Non-zero total effect indices associated with **sea ice parameters** are seen for several QOIs

GSA results: sensitivity indices


¹ Measure effect of individual parameters acting alone

² Measure variance that remains after learning values of every variable except z_i


³ Measure contribution of interaction between parameter subset u on the variance of a QOI




Main Effects¹



Sobol Indices²



Total Effects³



Sea ice
params

Atmosphere
params

Ocean
params

z_5 (clubb_c1) and z_6 (clubb_c8) combined effects are stronger than total effects for 5 insensitive parameters

- Figure above shows sensitivity indices for Sea Ice Volume QOI in spring
- Atmospheric parameters related to cloud parameterizations (CLUBB) are most important for all QOIs; z_6 (clubb_c8) has largest total effects
- Analysis also reveals significant parameter interactions between atmosphere parameters
- Non-zero total effect indices associated with sea ice parameters are seen for several QOIs
- Effects of ocean parameters and their interactions with each other/other parameters are negligible

GSA results: sensitivity indices


¹ Measure effect of individual parameters acting alone

² Measure variance that remains after learning values of every variable except z_i


³ Measure contribution of interaction between parameter subset u on the variance of a QOI




Main Effects¹



Sobol Indices²



Total Effects³



Sea ice
params

Atmosphere
params

Ocean
params

z_5 (clubb_c1) and z_6 (clubb_c8) combined effects are stronger than total effects for 5 insensitive parameters

- Figure above shows **sensitivity indices** for **Sea Ice Volume QOI** in **spring**
- Atmospheric parameters related to cloud parameterizations (CLUBB) are **most important** for all QOIs; z_6 (clubb_c8) has largest total effects
- Analysis also reveals **significant parameter interactions** between atmosphere parameters
- Non-zero total effect indices associated with **sea ice parameters** are seen for several QOIs
- Effects of **ocean parameters** and their interactions with each other/other parameters are **negligible**

Results are **qualitatively similar** to MOAT fully-coupled E3SMv0 study in [Urrego-Blanco *et al.*, 2019]

GSA results: sensitivity indices


¹ Measure effect of individual parameters acting alone

² Measure variance that remains after learning values of every variable except z_i


³ Measure contribution of interaction between parameter subset u on the variance of a QOI




Main Effects¹



Sobol Indices²



Total Effects³



Sea ice
params

Atmosphere
params


Ocean
params

z_5 (clubb_c1) and z_6 (clubb_c8) combined effects are stronger than total effects for 5 insensitive parameters

- Figure above shows sensitivity indices for Sea Ice Volume QOI in spring
- Atmospheric parameters related to cloud parameterizations (CLUBB) are most important for all QOIs; z_6 (clubb_c8) has largest total effects
- Analysis also reveals significant parameter interactions between atmosphere parameters
- Non-zero total effect indices associated with sea ice parameters are seen for several QOIs
- Effects of ocean parameters and their interactions with each other/other parameters are negligible

Although atmospheric parameters are most influential, they influence sea ice and ocean QOIs → analysis requires fully-coupled E3SM

GSA results: total effects indices – atmosphere parameters



Conclusions drawn from atmosphere parameters' total effects indices can be related to physical processes.

- SIE¹ QOI shows **strong response** to z_6 (clubb_c8) in fall
 - Increasing z_6 (clubb_c8) **brightens clouds**, resulting in Earth surface **cooling** [Larson, 2020]
 - Suggests that **cloud brightening** has potential to control **degree** to which **sea ice** is lost towards the end of the melting season
- Sensitivities in SIE & SIV² have strong **cyclical trends**
 - For z_4 (cldfrc_dp1) and z_7 (gamma_coeff), SIE¹ and SIV² **trend differently** – could reflect difference between **young, seasonal ice** and relatively stable **multi-year ice**
- Sensitivity of CLDLOW³ to clubb_c1 (z_5) in fall is **not as strong** as sensitivity of FLNS⁴
 - Results suggest while clubb_c1 (z_5) influences **cloud type**, it may not strongly influence the fraction of **general low cloud cover**

⋮

¹ Sea Ice Extent. ² Sea Ice Volume. ³ Low Cloud Overage below 700 hPa over Arctic. ⁴ Net longwave flux at surface over Arctic.



- Sensitivity analysis results can be used to calculate **normalized posterior mean of the main effect functions marginalized over one parameter at a time**, ± 2 standard deviations:

$$\mathbb{V}^*[Y]^{-\frac{1}{2}}(\mathbb{E}^*[\mathbb{E}[Y|z_i]] - \mathbb{E}^*[\mathbb{E}[Y]]) \pm 2\mathbb{V}^*[Y]^{-\frac{1}{2}}\mathbb{V}^*[\mathbb{E}[Y|z_i]]^{1/2}$$

* : expectation over posterior distribution of GP




Figure: marginalized main effects for surface temperature (TS) QOI. 95% CIs shown in gray.


Results can help **provide confidence** in the ULR E3SM and **guide model spin-ups**.

- Decreasing z_5 / z_7 and/or increasing z_4 / z_6 will **bring down** surface temperature
- Increasing z_6 (clubb_c8) is known to lead to **cloud brightening** and **cooling** the Earth's surface
- Low values of z_5 (clubb_c1) favor **insolation-reducing stratiform clouds** \Rightarrow **cooling**
- Curvature** in z_1 (ksno) and z_3 (dragio¹) match trends in [Urrego-Blanco *et al.*, 2016]
- Results are **consistent with manual tuning trends** observed

¹ Neutral ocean-ice drag



- Background and motivation
- Methodology
- Model tuning and spin-up
- Evaluation of the ultra-low resolution (ULR) E3SM
- Global sensitivity analysis results
- Summary






- We performed a **GSA** involving **9 parameters** and **6 QOIs** spanning **3 climate components** (atmosphere, ocean, sea ice) using the **ULR configuration** of the **fully-coupled E3SMv1**
 - **First GSA** using E3SMv1
 - A study of this scope is currently **intractable** using higher-resolution scientifically-validated configurations (e.g., 1° E3SM)
- A **spin-up** of the ULR E3SMv1 was performed to achieve an **equilibrium climate**
- ULR E3SM reproduced **large-scale patterns** in TOA radiation, precipitation, zonal mean temperature and zonal mean wind compared to observational data (CERES-EBAF, ERA-Interim) and the 1° E3SM
- **Main effect, total effect** and **Sobol indices** were calculated using a fast **Gaussian Process (GP) emulator** from 139 75-year runs of ULR E3SMv1 using the PyApprox software
- QOI-QOI and parameter-parameter interactions using sensitivity indices were able to **reconcile** relationships with several **well-known Arctic feedbacks**
- The **atmospheric parameters** related to cloud physics (CLUBB model in EAM) and their interactions had the largest impact on the Arctic climate state
 - Parameters were shown to affect **QOIs** from **3 different climate components**
- **Marginalized main effects functions** demonstrated that trends uncovered by this study are consistent with **manual spin-up** of ULR E3SMv1 and **physical processes** underlying the CLUBB parameterization



USE OUT-OF-THE-BOX
ULR E3SM FOR
SCIENTIFIC STUDIES



ULR E3SM IS A
PLAUSIBLE PHYSICS-
BASED *SURROGATE*
FOR UQ STUDIES

Significance of this work:

- Our study can serve as a **baseline** for and **guide** future studies with higher-resolution models, if/when it is **tractable** to repeat our GSA using higher-resolution E3SM
- Results can be used to show **number of samples** needed to get even moderate **accuracy** in a sensitivity analysis with variety of parameters → useful for predicting **computational budget** for future GSAs

Future work:

- Augment present study with higher-fidelity ensemble data (e.g., medium-low resolution 2.7° E3SM), towards a **multi-fidelity GSA**



ESSOAr
Earth and Space Science Open Archive

Advanced Search Search ESSOAr

ABOUT US SUBMIT AUTHOR DASHBOARD

Earth and Space Science Open Archive >

This preprint has been submitted to and is under consideration at Journal of Advances in Modeling Earth Systems (JAMES). ESSOAr is a venue for early communication or feedback before peer review. Data may be preliminary. Learn more about preprints

Preprint • Open Access • You are viewing the latest version by default [v2]

Global sensitivity analysis using the ultra-low resolution Energy Exascale Earth System Model

Authors

Irina Kalashnikova Tezaur , Kara Peterson , Amy Powell , John Jakeman , Erika Roesler

Published Online: Thu, 14 Oct 2021 | <https://doi.org/10.1002/essoar.10508267.2>

Download PDF Cite Tools Share

Abstract

For decades, the Arctic has been warming at least twice as fast as the rest of the globe. As a first step towards quantifying parametric uncertainty in Arctic feedbacks, we perform a variance-based global sensitivity analysis (GSA) using a fully-coupled, ultra-low resolution (ULR) configuration of version 1 of the Department of Energy's Energy Exascale Earth System Model (E3SMv1). The study randomly draws 139 realizations of ten model parameters spanning three E3SMv1 components (sea ice, atmosphere and ocean), which are used to generate 75 year long projections of future climate using a fixed pre-industrial forcing. We quantify the sensitivity of six Arctic-focused quantities of interest (QOIs) to these parameters using main effect, total effect and Sobol sensitivity indices computed with a Gaussian process emulator. A sensitivity index-based ranking of model parameters shows that the atmospheric parameters in the CLUBB (Cloud Layers Unified by Binormals) scheme have significant impact on sea ice status and the larger Arctic climate. We also use the Gaussian process emulator to predict the response of varying each variable when the impact of other parameters are averaged out. These results allow one to assess the non-linearity of a parameter's impact on a QOI and investigate the presence of local minima encountered during the spin-up tuning process. Our study confirms the necessity of performing global analyses involving fully-coupled climate models, and motivates follow-on investigations in which the ULR model is compared rigorously to higher resolution configurations to confirm its viability as a lower-cost surrogate in fully-coupled climate uncertainty analyses.

Pre-print available at:

<https://www.essoar.org/doi/10.1002/essoar.10508267.2> (paper is under review for publication in *JAMES*)

Other references:


- K. Peterson, A. Powell, I. Tezaur, E. Roesler, J. Nichol, M. Peterson, W. Davis, D. Stracuzzi, D. Bull. "Arctic Tipping Points Triggering Global Change LDRD Final Report", *Sand No. 2020-9932*. Sandia National Laboratories, Albuquerque, NM (2020) [and references therein].
- Code: <https://github.com/karapeterson/E3SM>
- Data: https://github.com/karapeterson/E3SM_ULR_GSA_Data

Thank you for your attention!



Start of Back-Up Slides


Goal: characterize important factors influencing **interannual variability** and **Arctic sea ice loss** in the coupled Earth system



<https://oceanbites.org/sea-ice-and-albedo-should-we-be-worried/>

Why sea ice?


- Sea ice plays an important role in **modulating Earth's climate**
 - Reflects solar radiation
 - Influences ocean circulation
- Sea ice loss **impacts other parts** of the Earth system¹
 - Impacts to mid-latitude weather, potentially increases in winter storms and drought
 - Disruption of Atlantic ocean circulation
 - Increased coastal erosion
- Loss of sea ice is expected to **encourage maritime and commercial activity**, potentially contributing to geopolitical conflict²
 - Potential for new trans-Atlantic shipping routes




Smith & Stephenson 2013

¹ Cohen *et al.*, 2018; Cvijanovic *et al.*, 2018. Sévellec *et al.*, 2017.


² Smith & Stephenson, 2013; "Climate Change and Security in the Arctic", Center for Climate & Security Report, Jan. 2021.



Ultra-low resolution (ULR)
atmosphere grid ($\approx 7.5^\circ$)



Medium-low resolution (MLR)
atmosphere grid ($\approx 2.7^\circ$)




Standard resolution
atmosphere grid ($\approx 1^\circ$)

GSA workflow with fully-coupled E3SM




- Perform **spin-up** of **ultra-low resolution (ULR)** model to reach equilibrium conditions of global climate state
- Select $P = 9$ **parameters** and **parameter ranges** from the sea ice, ocean and atmosphere components of the E3SM that play a significant role in model response based on previous literature
- Select $K = 6$ **QOIs**, including sea ice extent and surface air temperature averaged over the Arctic
- Use DAKOTA toolkit¹ to N **generate samples** from the parameter distributions
- Run 75-year simulations for each sample using the **ULR version of E3SM** with **pre-industrial control forcing**
 - For our study, $N = 202$ perturbed runs were attempted, which led to $M = 139$ successful 75-year runs



Above: ULR atmosphere grid ($\approx 7.5^\circ$)

Below: ULR ocean/sea ice grid (240km or $\approx 2.2^\circ$)



The **ULR configuration** enables sufficient ensemble generation for **full GSA** with the **fully-coupled E3SM**!



¹ <https://dakota.sandia.gov>

- ULR E3SM configuration is $\approx 100 \times$ less expensive to run than the “standard” 1° resolution E3SM
 - Using the ULR E3SM, our study took $\approx 1.00 \times 10^6$ CPU hours on Sandia’s Skybridge HPC cluster
 - The same study using the standard 1° resolution E3SM would require $\approx 1.14 \times 10^8$ CPU hours

Sensitivity analysis of individual Earth system components:

- “Uncertainty quantification and global sensitivity analysis of the Los Alamos sea ice model”, Urrego-Blanco *et al.*, *JGR Oceans*, 2016.
 - Sobol sensitivity analysis of 1° resolution LANL CICE model on 39 sea ice parameters
- “Parametric sensitivity and uncertainty quantification in the version 1 of E3SM atmosphere model based on short perturbed parameter ensemble simulations”, Qian *et al.*, *JGR Atmospheres*, 2018.
 - Used short (3-day) simulations of 1° resolution EAM to study sensitivity w.r.t. 18 atmosphere parameters
- “Antarctic ice shelf-ocean interactions in high-resolution, global simulations using the E3SM Part 2: Sensitivity studies and model tuning”, Asay-Davis *et al.*, *Ocean Sciences Meeting, AGU*, 2018.
 - Used $\approx 30\text{km}$ ($\approx 1^\circ$) resolution in Southern Ocean and Antarctic continental shelf to perform local sensitivity analysis of 12 land ice + ocean parameters using the Greens’ function method (GFM).

Sensitivity analysis of Earth system models for polar quantities of interest:

- “Emergent relationships among sea ice, longwave radiation, and the Beaufort high circulation exposed through parameter uncertainty analysis”, Urrego-Blanco *et al.*, *JGR Oceans*, 2019.
 - Used 1° resolution E3SMv0 with MOAT method, 5 parameters and 24 ensemble members

Our work **builds** on this research by performing the **first Global Sensitivity Analysis** with a **fully-coupled E3SMv1**



See talk by N. Urban,
MS15, Tues PM

Sensitivity analysis of individual Earth system components:

- “Uncertainty quantification and global sensitivity analysis of the Los Alamos sea ice model”, Urrego-Blanco *et al.*, JGR Oceans, 2016.
 - Sobol sensitivity analysis of 1° resolution LANL CICE model on 39 sea ice parameters
- “Parametric sensitivity and uncertainty quantification in the version 1 of E3SM atmosphere model based on short perturbed parameter ensemble simulations”, Qian *et al.*, JGR Atmospheres, 2018.
 - Used short (3-day) simulations of 1° resolution EAM to study sensitivity w.r.t. 18 atmosphere parameters
- “Antarctic ice shelf-ocean interactions in high-resolution, global simulations using the E3SM Part 2: Sensitivity studies and model tuning”, Asay-Davis *et al.*, Ocean Sciences Meeting, AGU, 2018.
 - Used $\approx 30\text{km}$ ($\approx 1^\circ$) resolution in Southern Ocean and Antarctic continental shelf to perform local sensitivity analysis of 12 land ice + ocean parameters using the Greens’ function method (GFM).




Sensitivity analysis of Earth system models for polar quantities of interest:



- “Emergent relationships among sea ice, longwave radiation, and the Beaufort high circulation exposed through parameter uncertainty analysis”, Urrego-Blanco *et al.*, JGR Oceans, 2019.
 - Used 1° resolution E3SMv0 with MOAT method, 5 parameters and 24 ensemble members

Our work **builds** on this research by performing the **first Global Sensitivity Analysis** with a **fully-coupled E3SMv1**



- At these resolutions we cannot resolve some processes
- Can we resolve large scale dynamics we are interested in?







Global surface temperature versus sea ice extent (left) and sea ice volume (right) trends. Historical ensemble trends are computed for the years 1979-2014 and overlapping 35-year pseudo-ensembles are created from the pre-industrial control simulation for the computed trends.

Factors controlling sea ice decline



- To predict when we will see an Ice-Free Arctic need to understand
 - Long-term decline due to external forcing (CO_2)
 - Superimposed year-to-year and decade-to-decade variability
- No consensus on
 - How much internal variability has influenced decline
 - Most important factors influencing internal variability
 - Recent papers have looked at
- Two recent papers looked at this:
 - Ding et al., Fingerprints of internal drivers of Arctic sea ice loss in observations and model simulations, Nature Geoscience 2019.
 - Screen and Deser, Pacific Ocean Variability Influences the Time of Emergence of Seasonally Ice-Free Arctic Ocean, GRL 2019.



Winter

	SIE	SIV	SST	TS	CLDLOW	FLNS
SIE	1.0	0.77	-0.90	-0.98	0.44	-0.039
SIV		1.0	-0.57	-0.86	-0.0545	0.38
SST			1.0	0.87	-0.67	0.28
TS				1.0	-0.30	-0.096
CLDLOW					1.0	-0.77
FLNS						1.0

Summer

	SIE	SIV	SST	TS	CLDLOW	FLNS
SIE	1.0	0.85	-0.90	-0.92	0.89	-0.87
SIV		1.0	-0.66	-0.73	0.66	-0.59
SST			1.0	0.99	-1.0	0.97
TS				1.0	-0.99	0.95
CLDLOW					1.0	-0.98
FLNS						1.0

Relationships between QOIs are generally consistent with expectations:

- Positive correlations between SIE¹ & SIV², SST³ & TS⁴
- Negative correlations between SIE/SIV & SST/TS
- Negative correlation between CLDLOW⁵ & FLNS⁶, especially in warmer seasons: a lot of low cloud cover ⇒ less net longwave radiation flux at the surface
- Negative correlation between CLDLOW and SST/TS across all 4 seasons
 - In winter, cloud coverage expected to increase surface temperature → not observed in our data, may be due to biases from runs without sea ice coverage
- Lack of correlation between SIE & FLNS and SIV & CLDLOW in winter is contrary to results obtained using higher resolutions of E3SM [Urrego-Blanco *et al.*, 2019]

¹ Sea Ice Extent. ² Sea Ice Volume. ³ Sea Surface Temperature averaged over 60-90° N. ⁴ Air Temperature averaged over 60-90° N. ⁵ Low Cloud Overage below 700 hPa averaged over 60-90° N. ⁶ Net longwave flux at surface over 60-90° N.

Effects associated with the 4 atmosphere parameters [Larson, 2020]:

- z_4 (**cldfrc_dp1**): CLUBB parameter which controls cumulus cloud-formation convective regimes in E3SM
- z_5 (**clubb_c1**): CLUBB parameter which controls the balance of cumulus versus stratocumulus clouds
 - Large positive values favor cumulus clouds, while small or negative values are associated with stratocumulus clouds
 - Stratocumulus clouds are believed to have planet-wide surface cooling effect, and Arctic cooling effects over most of the year [Eastman & Warren, 2010]
- z_6 (**clubb_c8**): CLUBB parameter developed to achieve radiative balance in atmospheric models
 - Increasing clubb_c8 brightens clouds, resulting in Earth surface cooling (brighter clouds reflect more incoming solar radiation)
- z_7 (**gamma_coeff**): tunable parameter in CLUBB shallow convection parameterization scheme that can brighten/dim clouds





Figure: cloud brightening produced micro-droplets that reflect more sunlight

<http://earthobservatory.nasa.gov/Features/Aerosols/page4.php>

GSA results: total effects indices – atmosphere parameters



*Atmosphere
parameters
influence
sea ice
QOIs*

*Figure: box
shows 25-75% CIs
for mean total
effects. Blue
dot indicates
mean.*

- For z_4 (cldfrc_dp1) and z_7 (gamma_coeff), SIE¹ and SIV² trend differently
 - Could reflect difference between relatively stable **multi-year ice** (measured by SIV) and **young, seasonal ice** (measured by SIE)
- SIE QOI shows **strong response** to z_6 (clubb_c8) in autumn
 - Increasing z_6 (clubb_c8) **brightens clouds**, resulting in Earth surface **cooling** [Larson, 2020]
 - Suggests that **cloud brightening** has potential to control **degree** to which **sea ice is lost** towards the end of the melting season

¹ Sea Ice Extent. ² Sea Ice Volume.

GSA results: total effects indices – atmosphere parameters




Figure: box shows 25-75% CIs for mean total effects. Blue dot indicates mean.

- CLDLOW¹ and FLNS² trend similarly for all but z_4 (cldfrc_dp1) parameter
- Sensitivity of CLDLOW to clubb_c1 (z_5) in autumn is not as strong as sensitivity of FLNS
 - clubb_c1 (z_5) parameter controls balance of cumulus vs. stratocumulus clouds [Larson, 2020]
 - Results suggest while clubb_c1 (z_5) influences cloud type, it may not strongly influence the fraction of general low cloud cover

¹ Low Cloud Overage below 700 hPa averaged over 60-90° N. ² Net longwave flux at surface over 60-90° N.



- [1] J. Urrego-Blanco, E. Hunke, N. Urban. "Emergent relationships among sea ice, longwave radiatino, and the Beaufort high circulation exposed through parameter uncertainty analysis". J. Geophys. Res. Oceans, 124, pp. 9572-9588, 2019. <https://doi.org/10.1029/2091JC014979>.
- [2] J. Urrego-Blanco, N. Urban, E. Hunke, A. Turner, N. Jeffery. "Uncertainty quantification and global sensitivity analysis of the Los Alamos sea ice model". J. Geophys. Res. Oceans, 121, pp. 2709-2732, 2016. doi: 10.1002/2015JC011558.
- [3] Y. Qian et al. "Parametric Sensitivity and Uncertainty Quantification in the Version 1 of E3SM Atmosphere Model Based on Short Perturbed Parameter Ensemble Simulations". J. Geophys. Res. Atmospheres, 123, pp. 13,046-13,073, 2018. <https://doi.org/10.1029/2018JD028927>.
- [4] P. Rasch et al. "An Overview of the Atmospheric Component of the Energy Exascale Earth System Model". J. Adv. Model. Earth Systems, 11, pp. 2377-2411, 2019. <https://doi.org/10.1029/1019MS001629>.
- [5] J.-C. Golaz et al. "The DOE E3SM Coupled Model Version 1: Overview and Evaluation at Standard Resolution". JAMES, 11, 2019. <https://doi.org/10.1029/2018MS001603>.
- [6] X. Asay-Davis, D. Comeau, J. Fyke, M. Hoffman, M. Petersen, S. Price, T. Ringler, L. Van Roekel, P. Wolfram. "Antarctic ice shelf-ocean interactions in high-resolution global simulations using the Energy Exascale Earth System Model (E3SM). Part 2: Sensitivity studies and model turning". 2018 Ocean Sciences Meeting, Portland, Oregon, Feb. 11-16, 2018.
- [7] S. Reckinger, M. Petersen, S. Reckinger. "A study of overflow simulations using MPAS-Ocean: vertical grids, resolution and viscosity". Ocean Modeling 96, pp. 291-313, 2015. <https://doi.org/10.1016/j.ocemod.2015.09.006>.



- Bathiany *et al.* Beyond bifurcation: using complex models to understand and predict abrupt climate change”, *DSCS* (2016).
- Cohen *et al.*, Arctic change and possible influence on mid-latitude climate and weather, US CLIVAR report (2018).
- Cohen, Pfeiffer and Francis, Warm Arctic episodes linked with increased frequency of extreme winter weather in the U.S, *Nature Communications* 9, article number 869, (2018).
- Cvijanovic *et al.* Future loss of Arctic sea-ice cover could drive a substantial decrease in California’s rainfall. *Nature Communications*, 8 (1), (2018).
- Duraisamy, *et al.* Turbulence Modeling in the Age of Data. *arXiv preprint arXiv:1804.00183* (2018).
- Enderlin, *et al.* An improved mass budget for the Greenland ice sheet, *Geophys. Res. Lett.*, 41, 866–872, (2014).
- Francis and Skific, evidence linking rapid Arctic warming to mid-latitude weather patterns. *Phil. Trans. R. Soc. A* 373: 20140170 (2015).
- Graeter *et al.* Ice Core Records of West Greenland Melt and Climate Forcing, *GRL*, 45, 3164-3172, (2018)
- Kinnard *et al.*, Reconstructed changes in Arctic sea ice over the past 1450 years, *Nature*, 479, 509-512 (2011).
- Koven *et al.*, Permafrost carbon-climate feedbacks accelerate global warming *PNAS*, volume 108, (2011).
- Lara *et al.*, Reduced arctic tundra productivity linked with landform and climate change interactions, *Scientific Reports*, volume 8, (2018).
- Lind, Ignvaldsen, and Furevik, Arctic warming hotspot in the northern Barents sea linked to declining sea-ice import, *Nature Climate Change*, 8 634-639 (2018).
- Notz and Stroeve, Observed Arctic sea-ice loss directly follows anthropogenic CO2 emission *Science* (2016).
- Olefeldt, D. *et al.* Circumpolar distribution and carbon storage of thermokarst landscapes. *Nat. Commun.* 7, 13043 (2016).
- Parazoo *et al.* Detecting the permafrost carbon feedback: talik formation and increased cold-season respiration as precursors to sink-to-source transitions *The Cryosphere*, 12, 123–144, (2018).
- Ricciuto, Sargsyan, Thornton, The Impact of Parametric Uncertainties on Biogeochemistry in the E3SM Land Model, *JAMES*, (2018).
- Schädel *et al.*, Potential carbon emissions dominated by carbon dioxide from thawed permafrost soils, *Nature Climate Change*, 6, (2016).
- Schuur *et al.*, Climate change and the permafrost carbon feedback, *Nature*, 520, (2015).
- Sévellec, Fedorov, Liu, Arctic sea-ice decline weakens Atlantic Meridional Overturning Circulation, *Nature Climate Change*, 7, 604-610 (2017).
- Smith and Stephenson, New trans-Arctic shipping routes navigable by mid-century, *PNAS* 110:13, 4871-4872 (2013).
- Strauss *et al.*, Deep Yedoma permafrost: A synthesis of depositional characteristics and carbon vulnerability, *Earth-Science Reviews*, (2017).
- Stroeve *et al.*, Trends in Arctic sea ice extent from CMIP5, CMIP3 and observations *GRL* (2012).
- van Angelen, *et al.*, Contemporary (1960–2012) Evolution of the Climate and Surface Mass Balance of the Greenland Ice Sheet, *Surv. Geophys.*, 35, 1155–1174, (2013).
- Vernon, *et al.* Surface mass balance model intercomparison for the Greenland ice sheet, *The Cryosphere* 7 599-614, (2013).